

Shrinkage estimators for prediction out-of-sample: Conditional performance

Nina Huber and Hannes Leeb

Department of Statistics, University of Vienna

Abstract

We find that, in a linear model, the James–Stein estimator, which dominates the maximum-likelihood estimator in terms of its in-sample prediction error, can perform poorly compared to the maximum-likelihood estimator in out-of-sample prediction. We give a detailed analysis of this phenomenon and discuss its implications. When evaluating the predictive performance of estimators, we treat the regressor matrix in the training data as fixed, i.e., we condition on the design variables. Our findings contrast those obtained by Baranchik (1973, *Ann. Stat.* 1:312–321) and, more recently, by Dicker (2012, arXiv:1102.2952) in an unconditional performance evaluation.

Keywords: James–Stein estimator, random matrix theory, random design

AMS subject classifications: 62M20, 62J07

1 Introduction

The problem of in-sample prediction, i.e., estimating the regression function at the observed design points, is arguably among the most extensively studied topics in regression analysis. But methods designed to perform well for in-sample prediction need not perform well for out-of-sample prediction, i.e., for estimating the regression function at a new point. We study the out-of-sample predictive performance of the James–Stein estimator, which dominates the maximum-likelihood estimator in the in-sample scenario; see Stein (1956) or the comprehensive monograph Judge and Bock (1978). We focus on the James–Stein estimator because of its conceptual importance (and because it is amenable to a detailed analytical analysis). The James–Stein estimator is the first method that was found to dominate maximum-likelihood through shrinkage, a discovery that helped to spark the development of many of the powerful estimation methods available today that rely on some sort of shrinkage through, e.g., regularization, model selection, or model averaging; see Leeb and Pötscher (2008) for a survey. In this paper, we find that the James–Stein estimator can perform poorly compared to the maximum-likelihood estimator in out-of-sample prediction, and we analyze and explain this phenomenon.

Consider the Gaussian linear regression model

$$Y = X\beta + u, \quad (1)$$

where X is a fixed $n \times p$ matrix of rank p , $\beta \in \mathbb{R}^p$, $n \geq p \geq 3$, and $u \sim N(0, \sigma^2 I_n)$. For simplicity, we focus on the known variance case and we assume that $\sigma^2 = 1$. Given an estimator $\tilde{\beta}$ for β , the corresponding in-sample prediction error, i.e., the mean squared error when estimating $X\beta$ by $X\tilde{\beta}$, will be denoted by $\rho_1(\tilde{\beta}, \beta, X)$ and is defined by

$$\rho_1(\tilde{\beta}, \beta, X) = \frac{1}{n} \mathbb{E} \left[(X\tilde{\beta} - X\beta)' (X\tilde{\beta} - X\beta) \right] = \mathbb{E} \left[(\tilde{\beta} - \beta)' \frac{X'X}{n} (\tilde{\beta} - \beta) \right]. \quad (2)$$

For out-of-sample prediction, consider a new set of explanatory variables, i.e., a p -vector x_0 , that is independent of Y , and hence also independent of $\tilde{\beta}$, and that satisfies $\mathbb{E}[x_0] = 0$ and $\mathbb{E}[x_0 x_0'] = \Sigma$, where Σ is positive definite and is regarded as a nuisance parameter. (In case of fixed x_0 , e.g., $x_0 = x_0^* \in \mathbb{R}^p$, we end up with the one-dimensional estimation target $x_0^{*'}\beta$, and it is well known that the maximum-likelihood estimator for $x_0^{*'}\beta$ is unique admissible minimax; see Lehmann and Casella (1998).) The out-of-sample prediction error is

the mean squared error when $x'_0\tilde{\beta}$ is used to predict $x'_0\beta$, where now the mean is taken with respect to both Y and x_0 . This error will be denoted by $\rho_2(\tilde{\beta}, \beta, X)$ and is defined by

$$\rho_2(\tilde{\beta}, \beta, X) = \mathbb{E} \left[(x'_0\tilde{\beta} - x'_0\beta)^2 \right] = \mathbb{E} \left[(\tilde{\beta} - \beta)' \Sigma (\tilde{\beta} - \beta) \right]. \quad (3)$$

(Of course, the out-of-sample prediction error $\rho_2(\tilde{\beta}, \beta, X)$ also depends on the matrix Σ , although this dependence is not explicitly shown in our notation.) We note that the existing results of Baranchik (1973) and Dicker (2012) consider prediction errors by assuming that X is random and by taking expectations as in (2) and (3) also with respect to X . We, on the other hand, compute prediction errors by treating X as fixed, i.e., we condition on the design. The expressions on the far right-hand sides of (2) and (3) differ in the matrices $X'X/n$ and Σ . If $X'X/n$ is very close to Σ , then the in-sample prediction error will be close to the out-of-sample prediction error. But if $X'X/n$ is not very close to Σ , then the in-sample predictive performance as measured by $\rho_1(\tilde{\beta}, \beta, X)$ can be markedly different from the out-of-sample predictive performance as measured by $\rho_2(\tilde{\beta}, \beta, X)$.

In this paper, we compare the maximum-likelihood estimator, the James–Stein estimator, and related shrinkage-type estimators by their performance as out-of-sample predictors. For in-sample prediction, i.e., in terms of the risk $\rho_1(\cdot, \cdot, X)$, it is well known that the James–Stein estimator dominates the maximum-likelihood estimator. But for out-of-sample prediction, i.e., in terms of $\rho_2(\cdot, \cdot, X)$, we find that the James–Stein estimator can perform quite poorly compared to the maximum-likelihood estimator; see Figure 1, relation (6), and also relation (8) in Theorem 4. (This finding contrasts a result of Baranchik (1973) as discussed at the end of Section 2 and after Theorem 3 in Section 3.) But we also find that such disappointing worst-case performance of the James–Stein estimator is atypical, in a certain sense; see relation (9) in Theorem 4.

For the case where Σ in (3) is known, estimators that dominate the maximum-likelihood estimator in terms of the risk (3) are well known, and we refer to Strawderman (2003) and the references given therein. The case where Σ is unknown but estimable is studied by Baranchik (1970), Berger and Bock (1976), Berger et al. (1977), and Copas (1983). For the challenging case where Σ is unknown and not estimable (in the sense that no further structural restrictions are imposed on Σ and that p/n is large, a scenario that we study in Section 3), we are not aware of further relevant existing results.

Explicit finite-sample formulae for the out-of-sample prediction errors of the estimators in question are derived in Section 2. In Section 3, we present approximations to the finite-sample quantities of interest; our approximations become accurate as $n \rightarrow \infty$, uniformly in the underlying parameters. Conclusions are drawn in Section 4, and the more technical derivations are collected in the appendices.

2 Explicit finite-sample results

Recall that the maximum-likelihood estimator of β is $\hat{\beta}_{ML} = (X'X)^{-1}X'Y \sim N(\beta, (X'X)^{-1})$. As pointed out by Stein (1956), James–Stein-type shrinkage estimators here correspond to estimators $\hat{\beta}(c)$ of β with $\hat{\beta}(c) = [1 - cp/(\hat{\beta}'_{ML}X'X\hat{\beta}_{ML})]\hat{\beta}_{ML}$, where $c \geq 0$ is a tuning-parameter (cf. Appendix A). In particular, the traditional James–Stein estimator corresponds to the estimator $\hat{\beta}(c)$ with $c = (p-2)/p$ and will be denoted by $\hat{\beta}_{JS}$; in the following, $\hat{\beta}_{JS}$ will also be called the James–Stein estimator (of β).

Proposition 1. *The in-sample prediction error and the out-of-sample prediction error of the James–Stein-type shrinkage estimator $\hat{\beta}(c)$ satisfy*

$$\rho_1(\hat{\beta}(c), \beta, X) = \rho_1(\hat{\beta}_{ML}, \beta, X) - \frac{1}{n} [2cp(p-2) - c^2p^2] \mathbb{E} \left[\frac{1}{\hat{\beta}'_{ML}X'X\hat{\beta}_{ML}} \right] \quad (4)$$

with $\rho_1(\hat{\beta}_{ML}, \beta, X) = p/n$, and

$$\begin{aligned} \rho_2(\hat{\beta}(c), \beta, X) &= \rho_2(\hat{\beta}_{ML}, \beta, X) - 2cp \text{trace}(\Sigma(X'X)^{-1}) \mathbb{E} \left[\frac{1}{\hat{\beta}'_{ML}X'X\hat{\beta}_{ML}} \right] \\ &\quad + (c^2p^2 + 4cp) \mathbb{E} \left[\frac{\hat{\beta}'_{ML}\Sigma\hat{\beta}_{ML}}{(\hat{\beta}'_{ML}X'X\hat{\beta}_{ML})^2} \right] \end{aligned} \quad (5)$$

with $\rho_2(\hat{\beta}_{ML}, \beta, X) = \text{trace}(\Sigma(X'X)^{-1})$, respectively.

The well-known formula on the right-hand side of (4) shows that the in-sample prediction error of $\hat{\beta}(c)$ equals the in-sample prediction error of $\hat{\beta}_{ML}$ minus the product of a positive expected value and a polynomial in c and p . In particular, $\rho_1(\hat{\beta}(c), \beta, X)$ is smaller than $\rho_1(\hat{\beta}_{ML}, \beta, X)$ if c satisfies $0 < c < 2(p-2)/p$ and is minimized for $c = (p-2)/p$, which is the tuning-parameter used by $\hat{\beta}_{JS}$. Moreover, it is easy to see that $\rho_1(\hat{\beta}(c), \beta, X)$ depends on β and X only through $\beta'(X'X/n)\beta$. Unlike the formula of $\rho_1(\hat{\beta}(c), \beta, X)$ in (4), display (5) shows that $\rho_2(\hat{\beta}(c), \beta, X)$ is obtained from $\rho_2(\hat{\beta}_{ML}, \beta, X)$ by subtracting a positive term and then adding another positive term, i.e., the second and the third term on the right-hand side of (5), that depend on c , on p , and on the unknown parameters in a more complicated fashion. By further inspection, we find that (5) depends on β and X through $\beta'(X'X/n)\beta$ and through $\beta'\Sigma\beta$ (which can be viewed as a kind of signal-to-noise ratio); for details, see Proposition A.1.




Figure 1: The solid curves in black and gray show finite sample relative prediction errors as a function of $\beta'\Sigma\beta$, and are explained in the next paragraph. The dashed curve shows the approximation to the relative prediction error that is obtained from Theorem 2 in Section 3. The constant solid line at 1 is for reference.

Figure 1 exemplifies the relative out-of-sample prediction error of the James–Stein estimator and of the maximum-likelihood estimator, i.e., $\rho_2(\hat{\beta}_{JS}, \beta, X)/\rho_2(\hat{\beta}_{ML}, \beta, X)$, as a function of $\beta'\Sigma\beta$ for various configurations in parameter space. For the figure, we selected a scenario where $X'X/n$ is not very close to Σ , so that the in-sample and the out-of-sample predictive performance of estimators differ from each other (as noted in the discussion after (3)). In particular, we took $n = 200$, $p = 160$, $\Sigma = I_p$, and X was obtained by sampling i.i.d. standard normals. The solid curves show $\rho_2(\hat{\beta}_{JS}, \beta, X)/\rho_2(\hat{\beta}_{ML}, \beta, X)$ as a function of $\beta'\Sigma\beta$, for β parallel to various eigenvectors of $X'X/n$. Let w_i be the eigenvector corresponding to the eigenvalue ν_i of $X'X/n$, and assume that $\nu_1 \leq \dots \leq \nu_{160}$. Four solid black curves stay below 1 and appear to be ordered; starting from the top, these correspond to β parallel to w_{160} , w_{120} , w_{80} , and w_{40} . The fifth solid black curve, that exceeds 1, corresponds to β parallel to w_1 , i.e., the eigenvector of the smallest eigenvalue. This curve attains a maximum of 4.27 at 75.12 (which is off the chart) and then recedes back towards 1 as $\beta'\Sigma\beta \rightarrow \infty$. The gray curves are obtained in the same way but for eigenvectors corresponding to the remaining smallest 25% of eigenvalues. The curves in Figure 1 differ dramatically depending on whether they correspond to small eigenvalues like ν_1 on the one hand, and moderate-to-large eigenvalues like ν_{40} ,

ν_{80} , ν_{120} , and ν_{160} on the other. Repeating these computations with X replaced by a new independent sample, we obtained essentially the same results. And for other choices of p and n , we obtained results that are qualitatively similar, including maxima above 1 corresponding to small eigenvalues. This phenomenon becomes less pronounced as p/n decreases, and it disappears completely for very small values of p/n . The results in Section 3 entail that this is no surprise.

Figure 1 shows that the James–Stein estimator no longer dominates the maximum-likelihood estimator for out-of-sample prediction, in the sense that

$$\rho_2(\hat{\beta}_{JS}, \beta, X) > \rho_2(\hat{\beta}_{ML}, \beta, X) \quad (6)$$

for some $\beta \in \mathbb{R}^p$, if X is the design matrix used to generate the figure. (Indeed, the left-hand side of the preceding display exceeds the right-hand side by a factor of 4.27 for appropriately chosen β , as noted in the preceding paragraph.) This should be compared to the findings of Baranchik (1973): The results in that paper suggest that

$$\mathbb{E} \left[\rho_2(\hat{\beta}_{JS}, \beta, X) \right] \leq \mathbb{E} \left[\rho_2(\hat{\beta}_{ML}, \beta, X) \right] \quad (7)$$

for each $\beta \in \mathbb{R}^p$, with strict inequality for some β , if X is random with i.i.d. $N(0, \Sigma)$ -distributed rows, and where the expectation in (7) is taken with respect to X (see also Dicker (2012)). Comparing the preceding two displays, we see that for X fixed, cf. (6), $\hat{\beta}_{JS}$ can perform poorly for some $\beta \in \mathbb{R}^p$. But on average with respect to X , as considered in Baranchik (1973) and Dicker (2012), the relation in (7) suggests that $\hat{\beta}_{JS}$ performs well, irrespective of β . The performance of $\hat{\beta}_{JS}$ hence depends crucially on whether we condition on the design as in (6) or average with respect to the design distribution as in (7). We give a more detailed analysis and explanation of this phenomenon in the next section. Also note that the phenomenon in (6) and (7) is related to the ancillarity paradox of Brown (1990).

3 Asymptotic approximations

In this section, we provide approximations to quantities like $\rho_2(\hat{\beta}_{JS}, \beta, X)$ and $\sup_{\beta} \rho_2(\hat{\beta}_{JS}, \beta, X)$ for ‘typical’ design matrices X . Here, ‘typical’ means ‘in probability’ when the explanatory variables in the training period, i.e., the rows of X , are taken as realizations from the same distribution as those in the prediction period (i.e., x_0). Our approximations are uniform in the unknown parameters and become accurate as $n \rightarrow \infty$, where the dimension of the model considered at sample size n , i.e., p , is allowed to depend on sample size. Note that quantities like $\rho_2(\hat{\beta}(c), \beta, X)$ and $\rho_2(\hat{\beta}_{ML}, \beta, X)$ now become random variables through their dependence on X . We emphasize that our evaluation of performance is always taken conditional on X , and that the random design is used only to describe the behavior for ‘typical’ design matrices X . We also stress, with X random, that the expectations in (4) and (5) are now to be understood as conditional on X .

For each n and p under consideration ($n \geq p \geq 3$), we assume that the model (1) holds; that x_0 and X are independent of the error u in (1); and that the rows of X and also x_0 are i.i.d. with mean zero and positive definite covariance matrix Σ . In addition, we assume that X can be written as $X = V\Sigma^{1/2}$, where $\Sigma^{1/2}$ denotes a symmetric square root of Σ and where V is the $n \times p$ matrix obtained by taking the upper left block of a double infinite array $(V_{i,j})_{i \geq 1, j \geq 1}$ of i.i.d. random variables that have mean zero, variance one, and a finite fourth moment (cf. Bai and Silverstein (2010)). Finally, we also assume that the (marginal) distribution of the $V_{i,j}$ ’s is absolutely continuous with respect to Lebesgue measure. This assumption could be dropped altogether, but at the expense of longer and technically more involved proofs. Under these assumptions, we note that $X'X$ is invertible almost surely. We set $\rho_2(\hat{\beta}(c), \beta, X) = 0$ on the probability-zero event where $X'X$ is degenerate. Otherwise, the random variable $\rho_2(\hat{\beta}(c), \beta, X)$ is defined by the expression on the right-hand side of (5), where the expected values are to be understood as conditional on X . These conventions also cover $\rho_2(\hat{\beta}_{ML}, \beta, X)$ and $\rho_2(\hat{\beta}_{JS}, \beta, X)$ in view of $\hat{\beta}_{ML} = \hat{\beta}(0)$ and $\hat{\beta}_{JS} = \hat{\beta}((p-2)/p)$. The following two results provide simple asymptotic approximations to quantities like $\rho_2(\hat{\beta}_{JS}, \beta, X)$ as well as $\sup_{\beta \in \mathbb{R}^p} \rho_2(\hat{\beta}_{JS}, \beta, X)$.

Theorem 2. *Assume that $n \rightarrow \infty$, and that $p = p(n)$ is such that $p/n \rightarrow t \in [0, 1)$. Moreover, for each p , let β and Σ be a p -vector and a positive definite $p \times p$ matrix, respectively, so that $\beta'\Sigma\beta \rightarrow \delta^2 \in [0, \infty]$ as $n \rightarrow \infty$. Then $\rho_2(\hat{\beta}_{ML}, \beta, X) \rightarrow t/(1-t)$ and $\rho_2(\hat{\beta}(c), \beta, X) \rightarrow r(\delta^2, c, t)$ in probability, where*

$$r(\delta^2, c, t) = \frac{t}{1-t} \left(1 - c \frac{t}{t + \delta^2} \right)^2 + c^2 \frac{t^2 \delta^2}{(t + \delta^2)^2}$$

if δ^2 is finite, and where $r(\infty, c, t) = t/(1-t)$ otherwise (expressions like $t/(t+\delta^2)$ are to be interpreted as zero if t and δ^2 are both equal to zero). Convergence of $\rho_2(\hat{\beta}(c), \beta, X)$ is uniform in c over compact sets, in the sense that $\sup_{0 \leq c \leq C} |\rho_2(\hat{\beta}(c), \beta, X) - r(\delta^2, c, t)| = o_P(1)$ for any $C > 0$. Moreover, these statements continue to hold if $\rho_2(\hat{\beta}(c), \beta, X)$ is replaced by the estimator $r(\hat{\delta}^2, c, p/n)$ where $\hat{\delta}^2 = \max\{Y'Y/n - 1, 0\}$.

Theorem 3. Assume that $n \rightarrow \infty$, and that $p = p(n)$ is such that $p/n \rightarrow t \in [0, 1)$. Moreover, for each p , let Σ be a positive definite $p \times p$ matrix. Then $\sup_{\beta \in \mathbb{R}^p} \rho_2(\hat{\beta}(c), \beta, X) \rightarrow \sup_{\delta^2 \geq 0} R(\delta^2, c, t)$ in probability, where

$$R(\delta^2, c, t) = \frac{t}{1-t} \left(1 - c \frac{t}{t + \delta^2}\right)^2 + c^2 \frac{t^2 \delta^2}{(1 - \sqrt{t})^2 (t + \delta^2)^2}$$

(again, expressions like $t/(t+\delta^2)$ are to be interpreted as zero if t and δ^2 are both equal to zero). Convergence is uniform in c over compact sets, so that $\sup_{0 \leq c \leq C} |\sup_{\beta \in \mathbb{R}^p} \rho_2(\hat{\beta}(c), \beta, X) - \sup_{\delta^2 \geq 0} R(\delta^2, c, t)| = o_P(1)$ for any $C > 0$.

Remark. The quantities $r(\delta^2, c, t)$ and $R(\delta^2, c, t)$, as defined in Theorems 2 and 3, respectively, differ by a factor of $(1 - \sqrt{t})^2$ in the denominator of the last term. The proofs reveal that this factor is caused by the fact that $\beta'(X'X/n)\beta$ can differ considerably from its expectation if p/n is not small, because of the gap between the smallest eigenvalue of a large-dimensional random Wishart matrix (or Gram matrix) and the smallest eigenvalue of its expectation, a well-known phenomenon in the theory of random matrices; see, for example, Bai and Silverstein (2010). In particular, in the setting of Theorem 2 with $\delta^2 = 1$, $\beta'(X'X/n)\beta$ converges to one in probability, but $\inf_{b, b' : \Sigma b = 1} b'(X'X/n)b$ converges to $(1 - \sqrt{t})^2$.

The approximations to $\rho_2(\hat{\beta}(c), \beta, X)$ and $\rho_2(\hat{\beta}_{ML}, \beta, X)$ provided by Theorem 2, i.e., $r(\delta^2, c, t)$ and $t/(1-t)$, respectively, are such that $r(\delta^2, c, t) < t/(1-t)$ whenever $t > 0$, provided only that $0 < c \leq 2$ and $\delta^2 < \infty$; cf. the dashed line in Figure 1. The results of Dicker (2012) suggest approximations to $\mathbb{E}[\rho_2(\hat{\beta}(c), \beta, X)]$ and $\mathbb{E}[\rho_2(\hat{\beta}_{ML}, \beta, X)]$ that coincide with $r(\delta^2, c, t)$ and $t/(1-t)$, respectively.

Theorems 2 and 3 together with the attending remark also provide us with a more precise description of the phenomenon in (6) and (7), and with a better understanding of the underlying cause. More formally, we have the following result.

Theorem 4. Assume that $n \rightarrow \infty$, and that $p = p(n)$ is such that $p/n \rightarrow t \in [0, 1)$. Moreover, for each p , let Σ be a positive definite $p \times p$ matrix. If $t > 1/9$, then the out-of-sample prediction errors of the James–Stein estimator and of the maximum-likelihood estimator are such that

$$\mathbb{P} \left(\sup_{\beta \in \mathbb{R}^p} \rho_2(\hat{\beta}_{JS}, \beta, X) - \rho_2(\hat{\beta}_{ML}, \beta, X) > \epsilon \right) \rightarrow 1 \quad (8)$$

for some $\epsilon > 0$ (that is given explicitly in the proof). And if $t \leq 1/9$, then the expression on the left-hand side of (8) converges to zero as $n \rightarrow \infty$ for each $\epsilon > 0$. Finally, irrespective of the value of $t \in [0, 1)$, we have

$$\sup_{\beta \in \mathbb{R}^p} \mathbb{P} \left(\rho_2(\hat{\beta}_{JS}, \beta, X) - \rho_2(\hat{\beta}_{ML}, \beta, X) > \epsilon \right) \rightarrow 0 \quad (9)$$

for each $\epsilon > 0$.

More generally, consider tuning-parameters $c_n \geq 0$ that converge to a limit $c \in [0, \infty)$ as $n \rightarrow \infty$, and consider the expression on the left-hand side of (8) with $\hat{\beta}(c_n)$ replacing $\hat{\beta}_{JS}$. The resulting expression converges to one as $n \rightarrow \infty$ for some $\epsilon > 0$ in the case where $0 \leq c \leq 2$ and $t > [(c-2)/(c+2)]^2$, and in the case where $c > 2$ and $t > 0$. In all other cases, the resulting expression converges to zero for each $\epsilon > 0$. And (9) holds for each $\epsilon > 0$ with $\hat{\beta}(c_n)$ replacing $\hat{\beta}_{JS}$ if $c \leq 2$, irrespective of t .

Remark. In the setting of Theorem 4, it is easy to see that relation (8) holds uniformly over all pairs of n and p subject to $1/9 + \delta \leq p/n \leq 1 - \delta$; in addition, (9) holds uniformly over all such pairs with $p/n \leq 1 - \delta$, for each $\delta > 0$, subject to $n \geq p \geq 3$ (and also uniformly over all positive definite $p \times p$ matrices Σ). Similar statements also apply with $\hat{\beta}(c_n)$ replacing $\hat{\beta}_{JS}$, mutatis mutandis.

Through relations (8) and (9), Theorem 4 provides two complementing views on the worst-case performance of James–Stein-type shrinkage estimators. If the expression on the left-hand side of (8) is large, then

the James–Stein estimator $\hat{\beta}_{JS}$ is typically outperformed, from a worst-case perspective, by the maximum-likelihood estimator $\hat{\beta}_{ML}$ (whose out-of-sample prediction error is constant in β). Here, ‘typically’ means that for most realizations of the design matrix X there is a parameter β for which $\hat{\beta}_{JS}$ performs worse than $\hat{\beta}_{ML}$. By Theorem 4, we see that this occurs with probability approaching one in the statistically challenging case where $t > 1/9$ (while this occurs with probability approaching zero in the case where $t \leq 1/9$). On the other hand, if the expression on the left-hand side of (9) is small, then with high probability X is such that $\hat{\beta}_{JS}$ outperforms $\hat{\beta}_{ML}$, uniformly in β . In the setting of Theorem 4, the left hand-side of (9) always converges to zero.

If $\rho_2(\cdot, \cdot, X)$ is used as a risk-function, then (8) entails that $\hat{\beta}_{JS}$ is outperformed by $\hat{\beta}_{ML}$ in terms of worst-case risk, for most realizations of the design matrix X . This is a worst-case perspective, as is often adopted in frequentist statistical analyses. But the most unfavorable parameter β , for which $\rho_2(\hat{\beta}_{JS}, \beta, X)$ is maximized, heavily depends on X . In particular, the relation in (9) entails, for any fixed parameter β , that the probability, that X is such that β is unfavorable, is small.

4 Discussion

We have derived explicit finite sample formulae for the out-of-sample prediction error of the James–Stein estimator and of related James–Stein-type shrinkage estimators in a linear regression model with Gaussian errors and fixed design. In an example with a particular design matrix X , we have found that the James–Stein estimator no longer dominates the maximum-likelihood estimator. We have shown that this phenomenon generally occurs for most design matrices X if the ratio of the number of explanatory variables in the model (p) and the sample size (n) exceeds $1/9$, in the sense of statement (8) of Theorem 4. At the same time, we have also shown that the James–Stein estimator outperforms the maximum-likelihood estimator for most design matrices X , uniformly in the underlying parameters, in the sense of statement (9) of Theorem 4. Our findings suggest that the James–Stein estimator can perform poorly for prediction out-of-sample from a frequentist worst-case perspective. But our findings also suggest, in the setting considered here, that the worst-case performance does not properly reflect the performance in the typical case, and that the James–Stein estimator performs favorably compared to maximum-likelihood in the typical case, uniformly in the underlying parameters.

The phenomenon, that the James–Stein estimator dominates the maximum-likelihood estimator for in-sample prediction but can fail to do so for prediction out-of-sample, is linked to the fact that the eigenvalues of $X'X/n$ can differ from the eigenvalues of Σ ; see relations (2) and (3), as well as the remark following Theorem 3. We therefore expect that other estimators, that are designed to perform well for in-sample prediction, can exhibit similar phenomena when used for prediction out-of-sample. This includes various shrinkage estimators like estimators based on model selection, penalized maximum-likelihood, and other forms of regularization. Although beyond the scope of this paper, it would be particularly interesting to study bridge estimators (Frank and Friedman, 1993), in particular, the LASSO (Tibshirani, 1996) and ridge regression (Hörl and Kennard, 1970), as well as the Dantzig selector (Candes and Tao, 2007) with regards to their performance as predictors out-of-sample when p/n is not small.

A Technical details for Section 2

Recall that $\hat{\beta}_{ML} \sim N(\beta, (X'X)^{-1})$, and define Z and ζ as $Z = (X'X)^{1/2}\hat{\beta}_{ML}$ and $\zeta = (X'X)^{1/2}\beta$, respectively, where $(X'X)^{1/2}$ denotes a symmetric square root of $X'X$. Note that we have $Z \sim N(\zeta, I_p)$, that the unknown parameter $\beta \in \mathbb{R}^p$ corresponds to the unknown parameter $\zeta \in \mathbb{R}^p$ via $\beta = (X'X)^{-1/2}\zeta$, and that any estimator $\hat{\beta}$ for β corresponds to an estimator $\hat{\zeta}$ for ζ via the relation $\hat{\beta} = (X'X)^{-1/2}\hat{\zeta}$, and vice versa. In the Gaussian location model $Z \sim N(\zeta, I_p)$ with $\zeta \in \mathbb{R}^p$, $p \geq 3$, consider the James–Stein-type shrinkage estimator $\hat{\zeta}(c) = (1 - cp/Z'Z)Z$ with tuning parameter $c \geq 0$. The estimator $\hat{\zeta}(c)$ for ζ corresponds to the estimator $(X'X)^{-1/2}\hat{\zeta}(c)$ for β , and it is elementary to verify that $(X'X)^{-1/2}\hat{\zeta}(c)$ equals $\hat{\beta}(c)$.

Proof of Proposition 1: Define Z , ζ and $\hat{\zeta}(c)$ as in the preceding paragraph. For the in-sample prediction error of the maximum-likelihood estimator, note that $\rho_1(\hat{\beta}_{ML}, \beta, X) = (1/n) \mathbb{E}[(Z - \zeta)'(Z - \zeta)] = p/n$. For relation (4), recall that $\hat{\zeta}(c)$ satisfies $\mathbb{E}[(\hat{\zeta}(c) - \zeta)'(\hat{\zeta}(c) - \zeta)] = p - [2cp(p - 2) - c^2p^2] \mathbb{E}[1/Z'Z]$; cf. James and Stein (1961). To verify that this equality is equivalent to (4), first note that the expression on the

left-hand side of this equality satisfies $\mathbb{E}[(\hat{\zeta}(c) - \zeta)'(\hat{\zeta}(c) - \zeta)] = n\rho_1(\hat{\beta}(c), \beta, X)$ (by plugging the definitions of $\hat{\zeta}(c)$, Z and ζ into the left-hand side, and by recalling the definitions of $\hat{\beta}(c)$ and $\rho_1(\hat{\beta}(c), \beta, X)$). And, in a similar fashion, the right-hand side of that equality is equal to the expression on the right-hand side of (4), multiplied by n .

For (5), we first use (3) together with the definition of $\hat{\beta}(c)$ to obtain that

$$\rho_2(\hat{\beta}(c), \beta, X) = \rho_2(\hat{\beta}_{ML}, \beta, X) - 2cp \mathbb{E} \left[\frac{\hat{\beta}'_{ML} \Sigma (\hat{\beta}_{ML} - \beta)}{\hat{\beta}'_{ML} X' X \hat{\beta}_{ML}} \right] + c^2 p^2 \mathbb{E} \left[\frac{\hat{\beta}'_{ML} \Sigma \hat{\beta}_{ML}}{(\hat{\beta}'_{ML} X' X \hat{\beta}_{ML})^2} \right]. \quad (10)$$

The expected value in the second term on the right-hand side of (10) can be written as

$$\mathbb{E} \left[\frac{Z' T (Z - \zeta)}{Z' Z} \right] = \sum_{i=1}^p \mathbb{E} \left[\frac{Z_i T_{i,i} (Z_i - \zeta_i)}{\sum_{k=1}^p Z_k^2} \right] + \sum_{\substack{i,j=1 \\ i \neq j}}^p \mathbb{E} \left[\frac{Z_j T_{j,i} (Z_i - \zeta_i)}{\sum_{k=1}^p Z_k^2} \right], \quad (11)$$

where Z and ζ have been defined earlier, and where $T = (X'X)^{-1/2} \Sigma (X'X)^{-1/2}$. For each of the terms in the first sum on the right-hand side of (11), we obtain that

$$\mathbb{E} \left[\frac{Z_i T_{i,i} (Z_i - \zeta_i)}{\sum_{k=1}^p Z_k^2} \right] = \mathbb{E} \left[\frac{T_{i,i}}{\sum_{k=1}^p Z_k^2} \right] - 2 \mathbb{E} \left[\frac{T_{i,i} Z_i^2}{(\sum_{k=1}^p Z_k^2)^2} \right]$$

upon using Stein's Lemma conditional on the Z_k 's with $k \neq i$. And each of the terms in the second (double) sum on the right-hand side of (11) can be written as

$$\mathbb{E} \left[\frac{Z_j T_{j,i} (Z_i - \zeta_i)}{\sum_{k=1}^p Z_k^2} \right] = -2 \mathbb{E} \left[\frac{T_{j,i} Z_i Z_j}{(\sum_{k=1}^p Z_k^2)^2} \right],$$

again by using Stein's Lemma conditional on the Z_k 's with $k \neq i$. Using the preceding two equalities, we can write the right-hand side of (11) as

$$\text{trace}(T) \mathbb{E} \left[\frac{1}{Z' Z} \right] - 2 \mathbb{E} \left[\frac{Z' T Z}{(Z' Z)^2} \right] = \text{trace}(\Sigma (X'X)^{-1}) \mathbb{E} \left[\frac{1}{\hat{\beta}'_{ML} X' X \hat{\beta}_{ML}} \right] - 2 \mathbb{E} \left[\frac{\hat{\beta}'_{ML} \Sigma \hat{\beta}_{ML}}{(\hat{\beta}'_{ML} X' X \hat{\beta}_{ML})^2} \right],$$

where the equality is obtained by using the definitions of T and Z . Recall that (11) or, equivalently, the right-hand side of the preceding display, is equal to the expected value in the second term on the right-hand side of (10). Plugging this into (10) and simplifying, we obtain (5). Furthermore, $\rho_2(\hat{\beta}_{ML}, \beta, X) = \text{trace}(\Sigma \mathbb{E}[(\hat{\beta}_{ML} - \beta)(\hat{\beta}_{ML} - \beta)']) = \text{trace}(\Sigma (X'X)^{-1})$. $\square$

The next result shows how (4) and (5) depend on β and on X ; moreover, that result can also be used to compute these expressions numerically.

Proposition A.1. *Assume that the linear model in (1) holds and that Σ is a positive definite $p \times p$ matrix. Then the expected values in (4) and (5) can be written as*

$$\begin{aligned} \mathbb{E} \left[\frac{1}{\hat{\beta}'_{ML} X' X \hat{\beta}_{ML}} \right] &= \mathbb{E} \left[\frac{1}{\chi_p^2(\beta' X' X \beta)} \right], \\ \mathbb{E} \left[\frac{\hat{\beta}'_{ML} \Sigma \hat{\beta}_{ML}}{(\hat{\beta}'_{ML} X' X \hat{\beta}_{ML})^2} \right] &= \mathbb{E} \left[\frac{\text{trace}(\Sigma (X'X)^{-1})}{(\chi_{p+2}^2(\beta' X' X \beta))^2} \right] + \mathbb{E} \left[\frac{\beta' \Sigma \beta}{(\chi_{p+4}^2(\beta' X' X \beta))^2} \right], \end{aligned}$$

where, for each $k \geq 1$ and each $\lambda \geq 0$, the symbol $\chi_k^2(\lambda)$ denotes a random variable that is chi-square distributed with k degrees of freedom and non-centrality parameter λ .

Proof. The first equality follows upon recalling that $X\hat{\beta}_{ML}$ follows a Gaussian distribution with mean $X\beta$ and covariance matrix $X(X'X)^{-1}X'$. The second equality is obtained by applying Corollary 2 from the appendix of Bock (1975). $\square$

B Technical details for Section 3

We start with some auxiliary results, and then prove the results from Section 3.

Lemma B.1. *For $t \geq 0$, $a = (1 - \sqrt{t})^2$ and $b = (1 + \sqrt{t})^2$, we have*

$$\int_a^b \frac{1}{x^2} \sqrt{(b-x)(x-a)} dx = 2\pi \frac{\min\{1, t\}}{|1-t|},$$

where the expression on the right-hand side of the preceding display is to be interpreted as ∞ in case $t = 1$.

Proof. The case $t = 0$ is trivial. In the case where $t > 0$, the integral of interest can be written as

$$\begin{aligned} \int_a^b \frac{1}{x^2} \sqrt{(b-x)(x-a)} dx &= 2(b-a)^2 \int_0^\infty \frac{z^2}{(az^2+b)^2(z^2+1)} dz \\ &= 2(b-a)^2 \int_0^\infty \left[\frac{a}{(b-a)^2} \frac{1}{(az^2+b)} + \frac{b}{b-a} \frac{1}{(az^2+b)^2} - \frac{1}{(b-a)^2} \frac{1}{z^2+1} \right] dz, \end{aligned}$$

where the first equality is obtained by the substitution $z = \sqrt{(b-x)/(x-a)}$, and the second equality follows from a partial fraction decomposition. Recall that $\int_0^w (u^2+1)^{-1} du = \arctan(w)$ and that $2k \int_0^w (u^2+1)^{-(1+k)} du = w(w^2+1)^{-k} + (2k-1) \int_0^w (u^2+1)^{-k} du$ whenever $k > 0$. With this, and for the case where $t \neq 1$, it follows that the integral on the far right-hand side of the preceding display reduces to

$$2(b-a)^2 \left[\frac{a}{(b-a)^2} \frac{\pi}{2\sqrt{a}\sqrt{b}} + \frac{b}{b-a} \frac{\pi}{4\sqrt{a}\sqrt{b^3}} - \frac{1}{(b-a)^2} \frac{\pi}{2} \right] = \frac{\pi}{2\sqrt{a}\sqrt{b}} (\sqrt{b} - \sqrt{a})^2.$$

Now note that $\sqrt{a} = 1 - \sqrt{t}$ in case $t < 1$, while $\sqrt{a} = \sqrt{t} - 1$ in case $t > 1$. This implies that the expression on the far right-hand side of the preceding display reduces to $2\pi \min\{1, t\}/|1-t|$, as required. And in case $t = 1$, we have $a = 0$ and $b = 4$, so that the integrand on the far right-hand side of the second-to-last display reduces to $[1 - 1/(z^2+1)]/16$ and hence integrates to ∞ . $\square$

Lemma B.2. *For the $n \times p$ matrix V as in Section 3, write $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p$ for the ordered eigenvalues of $V'V$. If n and p are such that $n \rightarrow \infty$ and $p/n \rightarrow t$ for some $t \in [0, 1]$, then*

$$\sum_{i=1}^p \frac{1}{\lambda_i} \xrightarrow{a.s.} \frac{t}{1-t},$$

where the limit is to be interpreted as ∞ in the case where $t = 1$ (and where the sum on the left-hand side is to be interpreted as ∞ whenever $\lambda_p = 0$). Moreover, we also have $\lambda_1/n \rightarrow (1 + \sqrt{t})^2$ and $\lambda_p/n \rightarrow (1 - \sqrt{t})^2$ almost surely.

Proof. Consider first the case where t satisfies $0 < t < 1$. Write $\nu_1 \geq \nu_2 \geq \dots \geq \nu_p$ for the ordered eigenvalues of $V'V/n$, and note that $\nu_i = \lambda_i/n$, $1 \leq i \leq p$. Write F_n for the empirical cumulative distribution function (c.d.f.) of the ν_i 's, i.e., for $x \in \mathbb{R}$, $F_n(x)$ is the fraction of eigenvalues of $V'V/n$ that do not exceed x , and note that F_n is a random c.d.f. as it depends on $V'V/n$. Under the maintained assumptions, F_n converges weakly to a non-random limit F , except on a set of probability zero; cf. Theorem 3.6 of Bai and Silverstein (2010). The limit c.d.f. F corresponds to the so-called Marchenko–Pastur distribution, which is supported by the interval $[a, b]$ with $a = (1 - \sqrt{t})^2$ and $b = (1 + \sqrt{t})^2$, and which has a density given by $f(x) = \sqrt{(x-a)(b-x)}/(2\pi tx)$ for $a \leq x \leq b$. For any bounded continuous function $g(\cdot)$, it follows that $\int g(x) F_n(dx) \rightarrow \int g(x) f(x) dx$ almost surely in view of the Portmanteau Theorem. And under the maintained assumptions, we also see that the smallest and the largest eigenvalue of $V'V/n$, i.e., $\nu_p = \lambda_p/n$ and $\nu_1 = \lambda_1/n$, converge almost surely to a and to b , respectively, whenever $0 < t \leq 1$; see Theorem 5.11 of Bai and Silverstein (2010). Recall that $a = (1 - \sqrt{t})^2$ is positive, and define a function $g(\cdot)$ as $g(x) = 1/x$ if $x \geq a/2$ and set $g(x) = 2/a$ otherwise. Now first note that $\int g(x) F_n(dx) \rightarrow \int g(x) f(x) dx$, except on a probability zero event, because $g(\cdot)$ is continuous and bounded. Second, we have that $\int g(x) f(x) dx = \int (1/x) f(x) dx$ because $g(x) = 1/x$ on the support of $f(\cdot)$. And third, observe that $\int g(x) F_n(dx)$ and $\int (1/x) F_n(dx)$ converge to the same limit,

except on a probability zero event, because ν_p converges to a almost surely. These three observations imply that

$$\int \frac{1}{x} F_n(dx) \xrightarrow{a.s.} \int \frac{1}{x} f(x) dx.$$

But the left-hand side in the preceding display can also be written as $\frac{1}{p} \sum_{i=1}^p \frac{1}{\nu_i} = \frac{n}{p} \sum_{i=1}^p \frac{1}{\lambda_i}$; and the right-hand side is equal to $1/(1-t)$ in view of Lemma B.1. It follows that $\sum_{i=1}^p 1/\lambda_i$ converges to $t/(1-t)$ almost surely, as required.

In the case where $t = 0$, choose an arbitrary number $\tilde{t}$ with $0 < \tilde{t} < 1$, and choose numbers $\tilde{p} \geq p$ that go to infinity with n such that $\tilde{p}/n \rightarrow \tilde{t}$ as $n \rightarrow \infty$. Write $\tilde{V}$ for the $n \times \tilde{p}$ matrix that forms the upper left block of the double array $(V_{i,j})_{i \geq 1, j \geq 1}$, and denote the eigenvalues of $\tilde{V}'\tilde{V}$ by $\tilde{\lambda}_1 \geq \dots \geq \tilde{\lambda}_{\tilde{p}}$. Because V is a sub-matrix of $\tilde{V}$, Cauchy's interlacing theorem entails that $\sum_{i=1}^p 1/\lambda_i \leq \sum_{i=1}^{\tilde{p}} 1/\tilde{\lambda}_i$, and also that $\tilde{\lambda}_{\tilde{p}} \leq \lambda_p \leq \lambda_1 \leq \tilde{\lambda}_1$. And from the case considered in the preceding paragraph, it follows that $\sum_{i=1}^{\tilde{p}} 1/\tilde{\lambda}_i \rightarrow \tilde{t}/(1-\tilde{t})$, that $\tilde{\lambda}_{\tilde{p}}/n \rightarrow (1-\sqrt{\tilde{t}})^2$, and that $\tilde{\lambda}_1/n \rightarrow (1+\sqrt{\tilde{t}})^2$, almost surely. Taken together, we obtain that $\limsup \sum_{i=1}^p 1/\lambda_i \leq \tilde{t}/(1-\tilde{t})$, that $(1-\sqrt{\tilde{t}})^2 \leq \liminf \lambda_p/n$, and that $\limsup \lambda_1/n \leq (1+\sqrt{\tilde{t}})^2$, almost surely. Letting $\tilde{t} \downarrow 0$ gives the desired result.

In the case where $t = 1$, we already know that $\lambda_1/n \rightarrow 4$ and $\lambda_p/n \rightarrow 0$ almost surely. Choose an arbitrary number $\tilde{t}$ with $0 < \tilde{t} < 1$, and let $\tilde{p} \leq p$ be such that $\tilde{p}/n \rightarrow \tilde{t}$. Now repeat the argument in the preceding paragraph, with the role of p and $\tilde{p}$ exchanged, to conclude that $\tilde{t}/(1-\tilde{t}) \leq \liminf \sum_{i=1}^p 1/\lambda_i$ almost surely. The result follows by letting $\tilde{t} \uparrow 1$. $\square$

Remark. The lemma continues to hold if $\sum_{i=1}^p 1/\lambda_i$ is replaced by $\sum_{i:\lambda_i > 0} 1/\lambda_i$. And the lemma can be adapted to cover also the case where $t \in (1, \infty]$.

Throughout the following, we use symbols like $\chi_k^2(\lambda)$ to denote a random variable that is chi-square distributed with $k \geq 1$ degrees of freedom and non-centrality parameter $\lambda \geq 0$. Note that $\chi_k^2(\lambda)$ has the same distribution as the sum of $k + 2J_\lambda$ i.i.d. central chi-square distributed random variables with one degree of freedom that are also independent of J_λ , where J_λ is Poisson with mean $\lambda/2$. In that sense, the law of $\chi_k^2(\lambda)$ can be viewed as a central chi-square distribution with random degrees of freedom equal to $k + 2J_\lambda$, and we will also denote this distribution by $\chi_{k+2J_\lambda}^2$. From this, it follows that $\chi_k^2(\lambda)$ is stochastically larger than $\chi_k^2(0)$, whence $\mathbb{E}[(1/\chi_k^2(\lambda))^m] \leq \mathbb{E}[(1/\chi_k^2(0))^m]$ for each $m \geq 1$ (where the expected values can be infinite). Also note that $\mathbb{E}[(1/\chi_k^2(0))^m] = \prod_{i=1}^m [1/(k-2i)]$ whenever k and m are positive integers satisfying $k > 2m$.

Lemma B.3. Fix an integer $m \geq 1$, and consider random variables $\chi_k^2(\lambda)$ with $k > 2m$. Then

$$\mathbb{E} \left[\left(\frac{k+\lambda}{\chi_k^2(\lambda)} \right)^m \right] \rightarrow 1$$

as $k + \lambda \rightarrow \infty$.

Proof. Fix $m \geq 1$. The mean and the variance of $\chi_k^2(\lambda)$ are given by $k + \lambda$ and $2(k + 2\lambda)$, respectively. Using this fact and Chebyshev's inequality, we see that $\chi_k^2(\lambda)/(k + \lambda) \rightarrow 1$, and also that $[(k + \lambda)/\chi_k^2(\lambda)]^m \rightarrow 1$, in probability as $k + \lambda \rightarrow \infty$. It remains to show, for integers k_a and reals $\lambda_a, a \geq 1$, satisfying $k_a > 2m$, $\lambda_a \geq 0$, and $k_a + \lambda_a \rightarrow \infty$ as $a \rightarrow \infty$, that the random variables $[(k_a + \lambda_a)/\chi_{k_a}^2(\lambda_a)]^m$ are uniformly integrable. In other words, for each fixed $\epsilon > 0$, we need to find a constant M so that

$$\mathbb{E} \left[\left(\frac{k_a + \lambda_a}{\chi_{k_a}^2(\lambda_a)} \right)^m \left\{ \left(\frac{k_a + \lambda_a}{\chi_{k_a}^2(\lambda_a)} \right)^m > M \right\} \right] < \epsilon \quad (12)$$

holds for each a , where we use symbols like $\{A\}$ to denote the indicator function of the event A . Set $U_a = [(k_a + \lambda_a)/\chi_{k_a}^2(\lambda_a)]^m$, note that the U_a 's are all integrable because $k_a > 2m$, and recall that $\chi_{k_a}^2(\lambda_a)$ is distributed as $\chi_{k_a+2J_a}^2(0)$ for an independent Poisson random variable J_a with mean $\lambda_a/2$. For later use, we also recall the Chernoff bound $\mathbb{P}(J_a < \lambda_a/4) \leq (e/2)^{-\lambda_a/4}$. To derive (12), i.e., to show uniform integrability of the U_a 's, we may assume that either $k_a \rightarrow \infty$ or $\lambda_a \rightarrow \infty$ as $a \rightarrow \infty$, by switching to subsequences if necessary (because $k_a + \lambda_a \rightarrow \infty$).

Assume that $k_a \rightarrow \infty$. Here, we first consider the (finitely many) a 's for which $k_a \leq 4m$. For these, we can find a constant M_1 so that (12) holds whenever $M \geq M_1$ because the U_a 's are integrable. It remains

to consider the a 's with $k_a > 4m$. For these a 's, we derive uniform integrability by showing that the second moments of the U_a 's are bounded. Using the law of iterated expectation, $\mathbb{E}[U_a^2]$ is given by the sum of

$$\mathbb{E} \left[\prod_{i=1}^{2m} \left(\frac{k_a + \lambda_a}{k_a - 2i + 2J_{\lambda_a}} \right) \{J_{\lambda_a} \geq \lambda_a/4\} \right] \quad (13)$$

and

$$\mathbb{E} \left[\prod_{i=1}^{2m} \left(\frac{k_a + \lambda_a}{k_a - 2i + 2J_{\lambda_a}} \right) \{J_{\lambda_a} < \lambda_a/4\} \right]. \quad (14)$$

The expression in (13) is bounded by $(4m+3)^{2m}$ because the i -th fraction in the integrand of (13) is bounded by $(k_a + \lambda_a)/(k_a - 2i + \lambda_a/2) \leq (k_a + \lambda_a)/(k_a - 4m + \lambda_a/2) \leq k_a/(k_a - 4m) + 2 \leq 4m + 1 + 2$ (the first inequality holds in view of $J_{\lambda_a} \geq \lambda_a/4$, and the last inequality holds because $k_a > 4m$). To bound the expression in (14), note that the i -th fraction in the integrand in (14) is bounded by $k_a/(k_a - 2i) + \lambda_a \leq 4m + 1 + \lambda_a$. Using this and the Chernoff's Poisson tail bound mentioned earlier, it follows that (14) is bounded by $(4m + 1 + \lambda_a)^{2m} (e/2)^{-\lambda_a/4}$. This upper bound is a continuous function of $\lambda_a \geq 0$ that converges to zero as $\lambda_a \rightarrow \infty$. Hence, the upper bound is itself bounded by a constant, uniformly in $\lambda_a \geq 0$, as required.

Now assume that $\lambda_a \rightarrow \infty$. We decompose the expression on the left-hand side of (12) into the sum of $\mathbb{E}[U_a \{U_a > M\} \{J_{\lambda_a} < \lambda_a/4\}]$ and $\mathbb{E}[U_a \{U_a > M\} \{J_{\lambda_a} \geq \lambda_a/4\}]$, and find M so that each of these two terms is bounded by $\epsilon/2$. To this end, note that $\chi_{k_a}^2(\lambda_a)$ is stochastically larger than $\chi_{k_a}^2(0)$, so that $\mathbb{E}[U_a \{U_a > M\} \{J_{\lambda_a} < \lambda_a/4\}]$ can be bounded by

$$\mathbb{E} \left[\left(\frac{k_a + \lambda_a}{\chi_{k_a}^2(0)} \right)^m \right] \mathbb{P}(J_{\lambda_a} < \lambda_a/4) \leq (2m + 1 + \lambda_a)^m (e/2)^{-\lambda_a/4}.$$

In the preceding display, the inequality is derived by writing the expected value as $\prod_{i=1}^m (k_a + \lambda_a)/(k_a - 2i)$, by noting that $k_a/(k_a - 2i) \leq 2m + 1$ because $k_a > 2m$, and upon using Chernoff's tail bound for the Poisson. Since $\lambda_a \rightarrow \infty$ here, the upper bound in the preceding display is smaller than $\epsilon/2$ for sufficiently large λ_a 's, e.g., $\lambda_a > \lambda_*$. And for the (finitely many) a 's for which $\lambda_a \leq \lambda_*$, we can find a constant M_2 so that $\mathbb{E}[U_a \{U_a > M\} \{J_{\lambda_a} < \lambda_a/4\}]$ is less than $\epsilon/2$ whenever $M \geq M_2$. Lastly, $\mathbb{E}[U_a \{U_a > M\} \{J_{\lambda_a} \geq \lambda_a/4\}]$ is bounded by

$$\begin{aligned} & \mathbb{E} \left[\left(\frac{k_a + \lambda_a}{\chi_{k_a + \lceil \lambda_a/2 \rceil}^2(0)} \right)^m \left\{ \left(\frac{k_a + \lambda_a}{\chi_{k_a + \lceil \lambda_a/2 \rceil}^2(0)} \right)^m > M \right\} \right] \\ & \leq 2^m \mathbb{E} \left[\left(\frac{k_a + \lceil \lambda_a/2 \rceil}{\chi_{k_a + \lceil \lambda_a/2 \rceil}^2(0)} \right)^m \left\{ \left(\frac{k_a + \lceil \lambda_a/2 \rceil}{\chi_{k_a + \lceil \lambda_a/2 \rceil}^2(0)} \right)^m > 2^{-m} M \right\} \right], \end{aligned}$$

where $\lceil x \rceil$ denotes the smallest integer not smaller than x , and where the inequality is based on the observation that $k_a + \lambda_a = 2(k_a/2 + \lambda_a/2) < 2(k_a + \lceil \lambda_a/2 \rceil)$. Set $\tilde{k}_a = k_a + \lceil \lambda_a/2 \rceil$ and set $\tilde{\lambda}_a = 0$. To find an $M_3 \geq M_2$ such that the expression on the right-hand side of the preceding display is less than $\epsilon/2$ whenever $M \geq M_3$, it suffices to show that the random variables $[(\tilde{k}_a + \tilde{\lambda}_a)/\chi_{\tilde{k}_a}^2(\tilde{\lambda}_a)]^m$ are uniformly integrable. Since $\tilde{k}_a \rightarrow \infty$ as $a \rightarrow \infty$, this has already been established in the second paragraph of the proof. $\square$

Lemma B.4. Consider the $n \times p$ matrix V as in Section 3, and let w be a unit-vector in $\mathbb{R}^p$ ($n \geq p \geq 3$). Then $\mathbb{E}[wV'Vw/n] = 1$ and $\text{Var}[wV'Vw/n] \rightarrow 0$ as $n \rightarrow \infty$, irrespective of the behavior of p ; in particular, we have $wV'Vw/n \rightarrow 1$ in probability.

Proof. Setting $W_i^{(n)} = (\sum_{j=1}^p V_{i,j} w_j)^2$, we see that $wV'Vw/n$ can be written as $(1/n) \sum_{i=1}^n W_i^{(n)}$, i.e., as the average of n i.i.d. random variables which have mean $\sum_{j=1}^p w_j^2 = 1$. And the variance of $W_i^{(n)}$ is given

by

$$\begin{aligned}
\sum_{a,b,c,d=1}^p w_a w_b w_c w_d \mathbb{E}[V_{i,a} V_{i,b} V_{i,c} V_{i,d}] - 1 &= \mathbb{E}[V_{1,1}^4] \sum_{a=1}^p w_a^4 + 3 \sum_{\substack{a,b=1 \\ a \neq b}}^p w_a^2 w_b^2 - 1 \\
&= (\mathbb{E}[V_{1,1}^4] - 1) \sum_{a=1}^p w_a^4 + 2 \sum_{\substack{a,b=1 \\ a \neq b}}^p w_a^2 w_b^2 \leq \mathbb{E}[V_{1,1}^4] + 1.
\end{aligned}$$

In the preceding display, the first equality is based on the fact that $\mathbb{E}[V_{i,a} V_{i,b} V_{i,c} V_{i,d}] = 0$ whenever one of the indices a, b, c, d differs from the others, while the second equality and the inequality are derived from the facts that $\sum_{j=1}^p w_j^2 = 1$ and that $1 \leq \mathbb{E}[V_{1,1}^4]$. We hence can bound $\text{Var}[w'V'Vw/n]$ by $(\mathbb{E}[V_{1,1}^4] + 1)/n$. $\square$

Lemma B.5. *Assume that Theorem 2 applies. If $t + \delta^2 > 0$, then*

$$\frac{p}{p + \beta' X' X \beta} \longrightarrow \frac{t}{t + \delta^2}, \quad \beta' \Sigma \beta \frac{p}{p + \beta' X' X \beta} \longrightarrow \frac{t \delta^2}{t + \delta^2},$$

in probability as $n \rightarrow \infty$; in case $\delta^2 = \infty$, the two limits are to be interpreted as 0 and as t , respectively. These statements continue to hold if p is replaced by $p + k$ in the numerators of the preceding display, for some fixed $k \in \mathbb{N}$.

Proof. Set $w = \Sigma^{1/2} \beta / (\beta' \Sigma \beta)^{1/2}$, note that $\|w\| = 1$, and that $\beta' X' X \beta / n$ can be written as $(\beta' \Sigma \beta) w' V' V w / n$. Using Lemma B.4, it follows that $\beta' X' X \beta / n \rightarrow \delta^2$ in probability. Because $\beta' X' X \beta / n$ and $\beta' \Sigma \beta$ both converge to δ^2 (the former in probability), and because $p/n \rightarrow t$, the continuous mapping theorem gives both limits in case $\delta^2 < \infty$, and the first limit in case $\delta^2 = \infty$. In the remaining case where $\delta^2 = \infty$, write the quantity of interest as $(p/n) / [(p/n) / \beta' \Sigma \beta + w' V' V w / n]$, which is easily seen to converge to t , as claimed. The last statement is trivial because p/n and $(p + k)/n$ converge to the same limit. $\square$

Proof of Theorem 2. For the maximum-likelihood estimator, we have $\rho_2(\hat{\beta}_{ML}, \beta, X) = \text{trace}(\Sigma(X'X)^{-1}) = \text{trace}((V'V)^{-1})$ almost surely, so that $\rho_2(\hat{\beta}_{ML}, \beta, X) \rightarrow t/(1 - t)$ in probability by Lemma B.2. For $\hat{\beta}(c)$, note that $\rho_2(\hat{\beta}(c), \beta, X)$ is given by (5) almost surely (where the expected values are to be understood as conditional on X , denoted by $\mathbb{E}[\cdot | X]$ in the following); using Proposition A.1 conditional on X , we can thus write $\rho_2(\hat{\beta}(c), \beta, X)$ also as

$$\begin{aligned}
&\text{trace}((V'V)^{-1}) \left(1 - 2c \mathbb{E} \left[\frac{p}{\chi_p^2(\beta' X' X \beta)} \middle| X \right] + (c^2 + 4c/p) \mathbb{E} \left[\frac{p^2}{(\chi_{p+2}^2(\beta' X' X \beta))^2} \middle| X \right] \right) \\
&+ (c^2 + 4c/p) \mathbb{E} \left[\frac{p^2 \beta' \Sigma \beta}{(\chi_{p+4}^2(\beta' X' X \beta))^2} \middle| X \right]
\end{aligned} \tag{15}$$

almost surely. Our first goal is to show that $\sup_{0 \leq c \leq C} |\rho_2(\hat{\beta}(c), \beta, X) - r(\delta^2, c, t)| = o_P(1)$.

In the case where $t + \delta^2 = 0$, note that $r(0, c, 0) = 0$, so that $\sup_{0 \leq c \leq C} |\rho_2(\hat{\beta}(c), \beta, X) - r(\delta^2, c, t)|$ is bounded from above by

$$\text{trace}((V'V)^{-1}) \left(1 + 2C \mathbb{E} \left[\frac{p}{\chi_p^2(0)} \right] + (C^2 + 4C/p) \mathbb{E} \left[\frac{p^2}{(\chi_{p+2}^2(0))^2} \right] \right) + (C^2 + 4C/p) \beta' \Sigma \beta \mathbb{E} \left[\frac{p^2}{(\chi_{p+4}^2(0))^2} \right]$$

because $\chi_{p+k}^2(\beta' X' X \beta)$ is stochastically larger than $\chi_p^2(0)$ for each $k \geq 0$. Because both $\text{trace}((V'V)^{-1})$ and $\beta' \Sigma \beta$ converge to zero (the former in probability) and the expected values are bounded by 3 for each $p \geq 3$, it follows that the expression in the preceding display converges to zero in probability.

In the case where $t + \delta^2 > 0$, we use the formula for $\rho_2(\hat{\beta}(c), \beta, X)$ in (15), the formula for $r(\delta^2, c, t)$, and the triangle inequality to bound $\sup_{0 \leq c \leq C} |\rho_2(\hat{\beta}(c), \beta, X) - r(\delta^2, c, t)|$ from above by

$$\begin{aligned} & \left| \text{trace}((V'V)^{-1} - \frac{t}{1-t}) + 2C \left| \text{trace}((V'V)^{-1} \mathbb{E} \left[\frac{p}{\chi_p^2(\beta'X'X\beta)} \right] X \right| - \frac{t^2}{(1-t)(t+\delta^2)} \right| \\ & + C^2 \left| \text{trace}((V'V)^{-1} \mathbb{E} \left[\frac{p^2}{(\chi_{p+2}^2(\beta'X'X\beta))^2} \right] X \right| - \frac{t^3}{(1-t)(t+\delta^2)^2} \right| \\ & + C^2 \left| \mathbb{E} \left[\frac{p^2 \beta' \Sigma \beta}{(\chi_{p+4}^2(\beta'X'X\beta))^2} \right] X \right| - \frac{t^2 \delta^2}{(t+\delta^2)^2} \right| \\ & + 4(C/p) \left(\text{trace}((V'V)^{-1}) \mathbb{E} \left[\frac{p^2}{(\chi_{p+2}^2(\beta'X'X\beta))^2} \right] X + \beta' \Sigma \beta \mathbb{E} \left[\frac{p^2}{(\chi_{p+4}^2(\beta'X'X\beta))^2} \right] X \right). \end{aligned}$$

We already know that the first term of the preceding display converges to zero in probability. For the remaining four terms, we note that either $p \rightarrow \infty$ (in case $t > 0$), or $\beta'X'X\beta \rightarrow \infty$ in probability (in case $\delta^2 > 0$ in view of Lemma B.4 because $n\beta'X'X\beta/n$ can be written as $n\beta'\Sigma\beta(w'V'Vw/n)$, where the unit-vector w is given by $\Sigma^{1/2}\beta/(\beta'\Sigma\beta)^{1/2}$). Using the assumption that $p \geq 3$, Lemma B.3 can be used to deal with the expected values in the preceding display. Together with the fact that $\text{trace}((V'V)^{-1}) \rightarrow t/(1-t)$ in probability, this shows that the sum of the four remaining terms converges to the same limit as

$$\begin{aligned} & 2C \frac{t}{1-t} \left| \frac{p}{p + \beta'X'X\beta} - \frac{t}{t + \delta^2} \right| + C^2 \frac{t}{1-t} \left| \frac{p^2}{(p + 2 + \beta'X'X\beta)^2} - \frac{t^2}{(t + \delta^2)^2} \right| \\ & + C^2 \left| \beta' \Sigma \beta \frac{p^2}{(p + 4 + \beta'X'X\beta)^2} - \frac{t^2 \delta^2}{(t + \delta^2)^2} \right| \\ & + 4(C/p) \left(\frac{t}{1-t} \frac{p^2}{(p + 2 + \beta'X'X\beta)^2} + \beta' \Sigma \beta \frac{p^2}{(p + 4 + \beta'X'X\beta)^2} \right). \end{aligned}$$

Lemma B.5 now entails that the expressions in the preceding display converge to zero in probability. (For the last term in the preceding display, we either have $p \rightarrow \infty$ if $t > 0$, or $t = 0$.)

Our second goal is to show that $\sup_{0 \leq c \leq C} |r(\hat{\delta}^2, c, p/n) - r(\delta^2, c, t)|$ also converges to zero in probability. We can bound the supremum in question by

$$\begin{aligned} & \left| \frac{p/n}{1-p/n} - \frac{t}{1-t} \right| + 2C \left| \frac{(p/n)^2}{(1-p/n)(p/n + \hat{\delta}^2)} - \frac{t^2}{(1-t)(t + \delta^2)} \right| \\ & + C^2 \left| \frac{(p/n)^3}{(1-p/n)(p/n + \hat{\delta}^2)^2} - \frac{t^3}{(1-t)(t + \delta^2)^2} \right| + C^2 \left| \hat{\delta}^2 \frac{(p/n)^2}{(p/n + \hat{\delta}^2)^2} - \delta^2 \frac{t^2}{(t + \delta^2)^2} \right|. \end{aligned}$$

The expression in the preceding display is the sum of four terms, where each term is of the form $|f(p/n, \hat{\delta}^2) - f(t, \delta^2)|$ for some function $f(\cdot, \cdot)$ which is continuous on $[0, 1) \times [0, \infty]$. Since $p/n \rightarrow t$ by assumption and $\hat{\delta}^2 \rightarrow \delta^2$ in probability (as is easy to see), the continuous mapping theorem entails that each of the four terms in the preceding display converges to zero in probability. $\square$

Proof of Theorem 3. On the almost-sure event where $X'X$ is invertible, $\rho_2(\hat{\beta}(c), \beta, X)$ is given by the formula (15). For invertible $X'X$ and for β such that $\beta'(X'X/n)\beta = d^2$, the expression in (15) depends on β only through $\beta'\Sigma\beta$ which is at most $d^2/\lambda_p(V'V/n)$, where $\lambda_p(\dots)$ denotes the smallest eigenvalue of the indicated matrix. [Indeed, we have $d^2 = \|(V'V/n)^{1/2}\Sigma^{1/2}\beta\|^2$ and $\beta'\Sigma\beta = \beta'\Sigma^{1/2}(V'V/n)^{1/2}[(V'V/n)^{-1}](V'V/n)^{1/2}\Sigma^{1/2}\beta \leq d^2/\lambda_p(V'V/n)$.] It follows that $\sup_{\beta \in \mathbb{R}^p} \rho_2(\hat{\beta}(c), \beta, X)$ can almost surely be written as the supremum of

$$\text{trace}((V'V)^{-1}) \left(1 - 2c \mathbb{E} \left[\frac{p}{\chi_p^2(nd^2)} \right] + (c^2 + 4c/p) \mathbb{E} \left[\frac{p^2}{(\chi_{p+2}^2(nd^2))^2} \right] \right) + \frac{c^2 + 4c/p}{\lambda_p(V'V/n)} \mathbb{E} \left[\frac{p^2 d^2}{(\chi_{p+4}^2(nd^2))^2} \right]$$

over $d^2 \geq 0$ or, equivalently, over $d \geq 0$. Write $R_*(d^2, c, n, p)$ for the random variable in the preceding display. The claim will follow if we can show that

$$\sup_{0 \leq c \leq C} \sup_{d \geq 0} |R_*(d^2, c, n, p) - R(d^2, c, t)|$$

converges to zero in probability. Now use the triangle inequality to bound the expression in the preceding display by the supremum of

$$\begin{aligned} & \left| \text{trace}((V'V)^{-1}) - \frac{t}{1-t} \right| + 2C \left| \text{trace}((V'V)^{-1}) \mathbb{E} \left[\frac{p}{\chi_p^2(nd^2)} \right] - \frac{t^2}{(1-t)(t+d^2)} \right| \\ & + C^2 \left| \text{trace}((V'V)^{-1}) \mathbb{E} \left[\frac{p^2}{(\chi_{p+2}^2(nd^2))^2} \right] - \frac{t^3}{(1-t)(t+d^2)^2} \right| \\ & + C^2 \left| \frac{1}{\lambda_p(V'V/n)} \mathbb{E} \left[\frac{p^2 d^2}{(\chi_{p+4}^2(nd^2))^2} \right] - \frac{t^2 d^2}{(1-\sqrt{t})^2(t+d^2)^2} \right| \\ & + 4C \text{trace}((V'V)^{-1}) \mathbb{E} \left[\frac{p}{(\chi_{p+2}^2(nd^2))^2} \right] + 4C \frac{1}{\lambda_p(V'V/n)} \mathbb{E} \left[\frac{p d^2}{(\chi_{p+4}^2(nd^2))^2} \right] \end{aligned}$$

over $d \geq 0$. The first term of the preceding display converges to zero in probability and each of the remaining five terms is of the form $|U_n f_n(d) - u f(d)|$ multiplied by a constant, where U_n is a random variable that converges to $u \in \mathbb{R}$ in probability (cf. Lemma B.2), and where $f_n(d)$ and $f(d)$ are functions of $d \in [0, \infty)$. For each of the remaining five terms, we now proceed as follows: To show that $\sup_{d \geq 0} |U_n f_n(d) - u f(d)| \rightarrow 0$ in probability, it suffices to show that $|U_n| \sup_{d \geq 0} |f_n(d) - f(d)| \rightarrow 0$ in probability (because $|U_n f_n(d) - u f(d)|$ is bounded by $|U_n| |f_n(d) - f(d)| + |f(d)| |U_n - u| \leq |U_n| |f_n(d) - f(d)| + |U_n - u|$, where the inequality follows upon noting that $f(d) \leq 1$). Let d_n , $n \geq 1$, be a sequence of maximizers (or near maximizers) of $|f_n - f|$. Because $[0, \infty]$ is compact, we may assume that $d_n \rightarrow \delta \in [0, \infty]$ (replacing the original sequence by a convergent subsequence if necessary). Moreover, it is easy to see that $\lim_{d \rightarrow \infty} f(d) = 0$ (for each of the five remaining terms in the preceding display). In particular, we can extend $f(\cdot)$ to a continuous function on $[0, \infty]$ by setting $f(\infty) = 0$. To complete the proof, we now show that either (a) $f_n(d_n) - f(\delta) \rightarrow 0$ as $n \rightarrow \infty$, or (b) $u = 0$ and $|f_n(d_n)|$ is bounded. In the case where $t + \delta^2 > 0$, we see that either $p \rightarrow \infty$ or $nd_n \rightarrow \infty$, and it is not difficult to conclude that case (a) occurs by using Lemma B.3 and the assumption that $p \geq 3$. And in the case where $t + \delta^2 = 0$, we see that case (b) occurs for the first, second, and fourth of the five remaining terms in the preceding display (by arguing as in the corresponding part in the proof of Theorem 2, mutatis mutandis), while again case (a) occurs for the third and the fifth as the expected values are bounded and d_n tends to zero. $\square$

Proof of Theorem 4. It suffices to show the statements for $\hat{\beta}(c_n)$. Set $R_* = \sup_{\delta^2 \geq 0} R(\delta^2, c, t)$ for $R(\cdot, \cdot, \cdot)$ as in Theorem 3, and note that $\sup_{\beta \in \mathbb{R}^p} \rho_2(\hat{\beta}(c_n), \beta, X) - \rho_2(\hat{\beta}_{ML}, \beta, X)$ converges to $R_* - t/(1-t)$ in probability by Theorems 2 and 3. For the first statement, assume either that $0 < c \leq 2$ and $t > [(c-2)/(c+2)]^2$ or that $c > 2$ and $t > 0$ hold (the case $c = 0$ can not occur there because $t < 1$). The statement now follows by observing, for such c and t , that $R_* - t/(1-t) > 0$, i.e., that $R(\delta^2, c, t) - t/(1-t) > 0$ for some $\delta^2 \geq 0$ (which is elementary but tedious to verify), and by setting ϵ equal to, say, $(R_* - t/(1-t))/2$. For the second statement, assume that $0 \leq c \leq 2$ and $t \leq [(c-2)/(c+2)]^2$, or that $c > 2$ and $t = 0$, and note, for such t and c , that $R_* - t/(1-t) \leq 0$, i.e., that $R(\delta^2, c, t) - t/(1-t) \leq 0$ for each $\delta^2 \geq 0$.

For the last statement, let $c \in [0, 2]$, and take $\epsilon > 0$. We need to show that $\mathbb{P}(\rho_2(\hat{\beta}(c_n), \beta, X) - \rho_2(\hat{\beta}_{ML}, \beta, X) > \epsilon)$ converges to zero for arbitrary sequences of parameters $\beta \in \mathbb{R}^p$ and Σ . Because the set $[0, \infty]$ is compact, we may assume that $\beta' \Sigma \beta$ converges to a limit $\delta^2 \in [0, \infty]$ (by considering convergent subsequences if necessary). It now follows from Theorem 2 that $\rho_2(\hat{\beta}(c_n), \beta, X)$ converges in probability to the limit $r(\delta^2, c, t)$ as is given in the theorem, while $\rho_2(\hat{\beta}_{ML}, \beta, X)$ converges in probability to $t/(1-t)$. The claim now follows from the fact that $r(\delta^2, c, t) \leq t/(1-t)$, irrespective of $\delta^2 \in [0, \infty]$ (which, again, is easy if somewhat tedious to verify). $\square$

References

- Z. Bai and J. W. Silverstein. *Spectral Analysis of Large Dimensional Random Matrices*. Springer, New York, 2nd edition, 2010.
- A. J. Baranchik. A family of minimax estimators of the mean of a multivariate normal distribution. *The Annals of Mathematical Statistics*, 41(2):642–645, 1970.
- A. J. Baranchik. Inadmissibility of maximum likelihood estimators in some multiple regression problems with three or more independent variables. *The Annals of Statistics*, 1(2):312–321, 1973.
- J. Berger and M. E. Bock. Combining independent normal mean estimation problems with unknown variances. *The Annals of Statistics*, 4(3):642–648, 1976.
- J. Berger, M. E. Bock, L. D. Brown, G. Casella, and L. Gleser. Minimax estimation of a normal mean vector for arbitrary quadratic loss and unknown covariance matrix. *The Annals of Statistics*, 5(4):763–771, 1977.
- M. E. Bock. Minimax estimators of the mean of a multivariate normal distribution. *The Annals of Statistics*, 3(1):209–218, 1975.
- L. D. Brown. An ancillarity paradox which appears in multiple linear regression. *The Annals of Statistics*, 18(2):471–493, 1990.
- E. Candes and T. Tao. The Dantzig selector: Statistical estimation when p is much larger than n . *The Annals of Statistics*, 35(6):2313–2351, 2007.
- J. B. Copas. Regression, prediction and shrinkage. *Journal of the Royal Statistical Society. Series B (Methodological)*, 45(3):311–354, 1983.
- L. Dicker. Dense signals, linear estimators, and out-of-sample prediction for high-dimensional linear models. *arXiv:1102.2952 [math.ST]*, 2012.
- I. E. Frank and J. H. Friedman. A statistical view of some chemometrics regression tools. *Technometrics*, 35(2):109–135, 1993.
- A. E. Hörl and R. W. Kennard. Ridge regression: Biased estimation for nonorthogonal problems. *Technometrics*, 12:55–67, 1970.
- W. James and C. Stein. Estimation with quadratic loss. *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability*, 1:361–379, 1961.
- G. G. Judge and M. E. Bock. *The statistical implications of pre-test and Stein-rule estimators in econometrics*. North-Holland Publishing Company, Amsterdam, 1978.
- H. Leeb and B. M. Pötscher. Model selection. In *Handbook of Financial Time Series*, pages 785–821. Springer, New York, 2008.
- E. L. Lehmann and G. Casella. *Theory of Point Estimation*. Springer, New York, 2nd edition, 1998.
- C. Stein. Inadmissibility of the usual estimator for the mean of a multivariate normal distribution. *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability*, 1:197–206, 1956.
- W. E. Strawderman. On minimax estimation of a normal mean vector for general quadratic loss. *Lecture Notes-Monograph Series*, 42:3–14, 2003.
- R. Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society. Series B (Methodological)*, 58(1):267–288, 1996.